



DECONTAMINATION

**ALTERNATIVE APPROACHES WHEN CLEANING &
DISINFECTION ARE NOT POSSIBLE
WORK INSTRUCTION & CHECKLIST**

Contents

1.Purpose 2

2.Actions 2

2.1 Required Materials & Personnel 4

2.2 Health & Safety Actions 4

🔗 Refer to: Avian influenza health and safety guidance for workplaces – Health New Zealand | Te Whatu Ora Error!

Bookmark not defined.

2.3 Burial Procedures Error! Bookmark not defined.

3.Appendices 5

Appendix A: Key Information 5

Appendix B: Options & Examples for Burial 5

Appendix C: Step-By-Step Instructions 6

Appendix E: Audit Checklist 9

Appendix F: Example SOP: Disposal by Composting (On-Site) (Farmer/operator use only) 9

Appendix F: Case Studies (Farmer/operator use only) 12

1.Purpose

This work instruction and checksheet outlines alternative process options, such as liming, stand-down, and fallowing, when cleaning and disinfection are not possible during a disease outbreak. The goal is to safely eliminate disease risk from a defined area to prevent persistence or spread and support the operation to resume safely.

Why This Matters:

In some situations, physical cleaning and disinfection may not be feasible due to site conditions, resource limitations, or contamination type. In these cases, alternative methods like liming and fallowing are critical to reduce disease risk and protect property, staff, and neighbouring operations. These approaches still require careful planning and monitoring to meet biosecurity obligations.

🔗 Please read this document in conjunction with the **Decontamination Guideline** and the **Disease Response Biosecurity Plan Guidebook** to ensure all procedures are followed correctly. Where a site already has a Standard Operating Procedure (SOP) for an alternative decontamination approach which meets the requirements in the Guideline, use the SOP.

In this document, “you” can refer to either an MPI person or a farmer/operator, depending on who is leading the response activities.

2.Actions

Before you choose the decontamination method, assess the risk level of each item or area. This helps match the right mitigation strategy to the type of contamination.

🔗 Use **Table 1** below to guide your decisions. It shows examples of contamination risks and the corresponding mitigation strategies.

Table 1. Risk Levels and Mitigation Strategies

Risk Level	Examples	Recommended Mitigation Strategies
Very High (5)	Carcasses, raw meat, contaminated eggs	Liming, stand-down period, full disinfection
High (4)	Slaughter equipment, used PPE, animal health equipment	Stand-down period, partial disinfection
Medium (3)	Vehicles, packaging, initial processed products	Fallowing, restricted access
Low (2)	Off-site vehicles, further processed products, biosecure feed	Routine cleaning, monitoring
Very Low (1)	Cooked products, diagnostic samples, general waste	Standard hygiene protocols

How This Fits Into The Plan

Knowing the contamination risks helps you:

- Select the right decontamination method (e.g. full disinfection vs. fallowing).
- Allocate appropriate gear and personnel.
- Develop a clear health and safety plan, including procedures for biosecure transport.

✂ **See Appendix A** for the key requirements an alternative decontamination plan must meet.

✂ **See Appendix B** for mitigation options and examples to help to choose the right approach.

✂ **See Appendix C** for step-by-step instructions on how to carry out alternative decontamination approaches.

Remember The Red Line

The *farm production area map* (Section A.2.1 of the Response Plan Template) shows the infected site. Treat the site like it has a border - the **red line**.

Inside the red line = contaminated.

Your job: **keep infection from crossing out.**

The plan should be based on:

- **Type of Contamination**
 - What's present (e.g. manure, blood, dust).
 - What disease organism is being targeted (e.g. HPAI virus, *Salmonella Enteritidis*).
- **Disease Spread and Cross-Contamination Risks**
 - Movement of animals, people, vehicles, and tools.
 - Traffic flow, barriers, and signage.
 - Nearby waterways or farms.
 - Weather and site conditions (e.g. temperature, wind, rain).
- **Sensitive Zones**
 - Areas needing exclusion or special handling (e.g. offices, feed storage, living quarters).

Ask yourself:

- ☒ How do I protect disinfected areas from being re-contaminated?
- ☒ How will I contain water run-off to prevent disease spread?

Work closely with contractors to ensure everyone understands the plan and follows the same protocols.

2.1 Required Materials & Personnel

These are the minimum requirements (liming only). You may need to scale up depending on the size of the job:

Category	Item
 Personal Protective Equipment (PPE)	• Coveralls. • Gloves. • Respirator. • Goggles.
 Safety Equipment	• Hydrated lime or quicklime. • Spreaders, shovels • Water source • Containment barriers

For stand-down and fallowing, signage and physical barriers such as fences and gates are required to ensure areas are separated and secured.

2.2 Health & Safety Actions

The on-site **written, Health & Safety plan** for decontamination activities at the infected site, needs to cover risks associated with alternative decontamination approaches during a disease incursion. If you're working with **contractors**, make sure you're familiar with and coordinate with their Health & Safety Plans too.

This ensures everyone is working safely, understands the risks, and follows the right procedures to prevent disease spread.

The plan should cover:

- Safe use of the equipment (e.g. applicators) and substances (e.g. lime).
- Entry and exit points (watch for cross-contamination), including disinfection steps before departure and after arrival in different zones (e.g. entering a shed) and on and off of the property.
- Risks from handling infected materials.
- Vehicle, equipment and personnel movement through high-risk zones (e.g. near sheds, water sources, or red line boundaries).
- PPE requirements.

Follow these rules:

- ☒ Conduct a health and safety briefing based on the site-specific plan.
- ☒ Keep accessways clear and safe for vehicles and machinery.
- ☒ Move from dirty to clean, never the other way around, to prevent cross-contamination.
- ☒ Ensure adequate lighting e.g. in sheds or when working at night.
- ☒ Wear correct PPE at all times.
- ☒ Stick to all biosecurity procedures, including recording and reporting any new risks.
- ☒ Handle materials carefully to prevent disease spread during decontamination.

 Refer to: [Avian influenza health and safety guidance for workplaces – Health New Zealand | Te Whatu Ora](#)

3. Appendices

Appendix A: Key Information

When physical cleaning and disinfection can't be done, alternative approaches like liming, stand-down, and fallowing must be planned and verified to reduce disease risk.

Plan your approach

- A clear plan ensures no steps are missed and contamination is contained.
- Match the method to the contamination type and risk level (see table 1 in main document).
- Use your farm map to define the red line, "the contaminated zone", and prevent spread beyond it.

Surface & site challenges

- Uneven terrain, porous materials, and hidden areas (e.g. ducts, pits, motors) reduce effectiveness.
- Seal or replace surfaces that can't be cleaned.
- Weather, water runoff, and nearby farms must be considered in your plan.

Water quality

- Poor water quality (e.g. High pH or hardness) can reduce disinfectant effectiveness.
- Check with your supplier if unsure as incorrect water use wastes time and leaves pathogens behind.

Self-assessment & verification

- Use torches, swabs, and checklists to confirm cleaning and mitigation steps.
- Clean before disinfecting, and verify before moving on.
- Contractors must follow the same protocols and health & safety plans.

Liming

- Use hydrated lime for surface areas and quicklime for burial pits (requires water slaking).
- Apply evenly, restrict access for 72 hours, and monitor at 30, 60, and 90 days.
- Lime is caustic – avoid contact and keep away from water bodies.
- Record type, quantity, location, personnel, and inspection results.

Stand-down

- Triggered when disease is absent or eradicated.
- Cleaning must be completed first.
- Applies to sites, vehicles, and tools – movement restrictions remain until verified.
- Mpi must assess and confirm recovery status.

Fallowing

- Restrict access for at least 30 days or as advised.
- Use signage and barriers to secure the area.
- Monitor for wildlife or unauthorised entry.
- Environmental sampling may be required.


Appendix B: Options & Examples for Burial

Liming



Liming involves applying lime to contaminated surfaces or burial pits to elevate pH and inactivate pathogens.

Stand-down



Stand-down involves cleaning followed by a period of non-use to allow any remaining pathogens to die off.

Fallowing



Fallowing refers to restricting access to an area over a specified period to contain pathogens and allow them to degrade naturally.

Appendix C: Step-By-Step Instructions

☒ Pre-Start Checklist

Action	Complete (Yes / No)
Electrical equipment made safe (if applicable)	
Bulk soil removed from all surfaces	
Sensitive equipment sprayed and wiped (MPI-approved)	
Sensitive equipment covered (e.g. Plastic bags)	
Vehicle wash/spray station set up at entry/exit	
Cleaning team briefed and PPE issued	
Waste disposal plan confirmed	
Water source and quality checked	
Detergents and disinfectants on hand (MPI-approved)	

Liming

Step-by-step instructions for applying lime to contaminated areas.

Materials and equipment:

- Hydrated lime or quicklime.
- PPE (coveralls, gloves, respirator, goggles).
- Spreaders, shovels.
- Water source.
- Containment barriers.

Safety Preparation:

- Full PPE required.
- Prepare site-specific health and safety plan.
- Store lime securely.

Lime Selection:

- Hydrated lime for surface use.
- Quicklime for burial pits (requires slaking with water).

Application:

- Apply lime evenly to affected areas as per site plan

Monitoring:

- Restrict access for 72 hrs.
- Inspect at 30, 60, 90 days.
- Monitor for recontamination.

Safety and Environment:

- Lime is caustic: avoid skin/eye contact
- Keep away from water bodies
- Dispose of unused lime safely

Documentation:

- Record lime type, quantity, application date/location, personnel involved, and inspection results
- Refer to verification checklist document

Stand-down

Step-by-step instructions for implementing a stand-down protocol.

First Step:

- Triggered when the disease is not detected or has been eradicated.
- Applies to both places and items (e.g., vehicles, equipment).

Cleaning and Disinfection:

- Must be completed before the stand-down period begins.
- Strategies should align with WOAHS standards where practicable.

Scope of Application:

- Includes commercial poultry sites, transport vehicles, tools, and equipment.
- Movement restrictions remain in place until stand-down is verified.

Verification and Assurance:

- Stand-down periods and cleaning/disinfection must be assessed in consultation with an MPI place-based manager.
- Verification ensures the site can be classified as recovered and disease-free.
- Robust stand-down periods serve as final assurance against ongoing transmission.

Fallowing

Step-by-step instructions for implementing fallowing procedures.

- Restrict access to fallowed areas for a minimum of 30 days or as advised by biosecurity authorities.
- Monitor the area for signs of wildlife or unauthorised entry.
- Maintain signage indicating restricted access and biosecurity status.

Conduct periodic environmental sampling if required.

Appendix E: Audit Checklist

GENERAL

Site name:	
Site address:	
Site contact name:	
Movement Control Permit/s number/s (as required) (for off-site):	

DOCUMENTATION

Document	Comment	Acceptable Y/N
Procedures (SOP).		
Contract related Statement of Work (where appropriate)		
Site Health and Safety Procedures		
Movement control permits, if required		

RECORDS

Records	Comment	Acceptable Y/N
Numbers of sites (Places) to be decontaminated		
Monitoring of the additional mitigation (liming, stand down and/or fallowing)		
Review scoring and outcomes and corrective actions		
Monitoring of cleaning and disinfection of vehicles (where appropriate)		

REALITY

Liming Procedure Checklist

Mitigation		Chemical concentration and/or Application Rate	Correct - √ or Needs Improve- X	Date and time
Liming Process for areas adjacent to sheds	Removal of any material which would interfere with liming efficacy			
	Liming			
	Treatment of material (example Pit Liming)			

Stand Down Protocol Checklist

Mitigation	Identification (area or place)	Signage (if applicable)	Date and time
Area restriction			
Monitor area			
Biosecurity status			
Environmental status			

Fallowing Procedure Checklist

Mitigation	Identification (area or place)	Signage (if applicable)	Date and time
Area restriction			
Monitor area			
Biosecurity status			
Environmental status			

Appendix F: Example SOP: Fallowing Procedure (Farmer/operator use only)

Purpose

To restrict access to contaminated areas for a defined period, allowing natural degradation of pathogens and ensuring biosecurity compliance.

Scope

This SOP applies to all areas designated for fallowing during a biosecurity response where cleaning and disinfection are not feasible.

Responsibilities

- Site Supervisor: Designate fallowing zones and ensure signage and barriers are in place.
 - Monitoring Team: Conduct regular inspections and environmental sampling.
-

Materials and Equipment

- Physical barriers (e.g. fences, gates).
 - Signage indicating restricted access.
 - PPE for monitoring personnel.
 - Environmental sampling kits.
-

Procedure

1. Site Preparation

- Identify and map fallowing zones, and uncontaminated areas that can be used.
- Install physical barriers to prevent access.
- Place clear signage indicating restricted access and biosecurity status.

2. Access Restriction

- Restrict access for a minimum of approximately 60-120 days or as advised by biosecurity authorities (see Decontamination Guidelines for specific details).
- Ensure no personnel, vehicles, or equipment enter the fallowed area.

3. Monitoring

- Conduct regular inspections for signs of wildlife or unauthorised entry.
- Maintain barrier integrity and signage visibility.

4. Environmental Sampling

- Perform periodic environmental sampling if required by biosecurity authorities.
 - Record and report findings to the site supervisor.
-

Records

- Fallowing zone map and designation records.
- Inspection logs.
- Environmental sampling results.
- Access restriction compliance reports.
- Corrective and preventative actions.

Appendix F: Case Studies (Farmer/operator use only)

When HPAI Hit: How I Fallowed My Small Free-Range Egg Farm and Survived¹

I run a small free-range egg farm with about 300 birds. We've got caravans and a few paddocks where the hens roam. When HPAI hit, it felt like the world stopped. Everything I'd built was suddenly at risk.

I looked at the options MPI gave us, full cleaning and disinfection, but the cost was way beyond what I could manage. The gear, the chemicals, the contractors... it just wasn't practical for a setup like mine. So I went with the only option that made sense: fallowing.

Euthanasing the birds was hard, but many of them were already sick so I felt I was stopping them suffering. Then, I scraped out every bit of manure from the caravans. It was hard, dirty work, but I got it done. I composted the birds, the manure, and the bedding on-site. Because the quantity was small, I didn't need a resource consent, which was a relief.

Then I limed the fields. Hydrated lime went down across the paddocks to knock back any virus that might still be hanging around. After that, I locked the place up, signs on the gates, no entry for anyone. The farm sat empty for twelve months, just me checking in now and then to make sure nothing had gone wrong.

I was lucky at the start as it was summer, hot and dry. The UV from the sun helped kill off anything left behind. That gave me some peace of mind.

I didn't even let the sheep back onto the paddocks. They had to go. I sold them off because I couldn't risk spreading the virus. That was another hit, but it had to be done.

The hardest part wasn't the work, it was the money. I had to go back to full-time work off-farm just to keep afloat. Everything went on the credit card, and I'm still paying it off. It's tough, but I did what I had to do to protect the farm and stop the spread.

If there's one thing I learned, it's that fallowing works when you've got no other choice. It's not easy, and it's not cheap, but it's doable. And sometimes, that's all you've got.

¹ This is a fictional scenario for illustrative purposes only.